

How do I know if SCIT will work for me?

If your doctor thinks that SCIT might help you, then you will be offered an appointment in a specialist allergy clinic. During the clinic appointment, a detailed history will be taken and skin prick tests will be performed to a range of pollens and other allergens. We will also ensure that, if you have asthma, it is well controlled so that you can have the treatment.

What if it doesn't work?

SCIT will not work for everybody. Pollinex Quattro leads to “very good” or “good” improvement in three out of four pollen allergic patients. It also reduces the use of hay fever medicines. If it doesn't work, patients usually have to rely on avoidance strategies and symptomatic treatment. Sometimes there are other allergies e.g. pet allergy or dust mite allergy that need to be sorted.

How can I contact you if I need to?

You will have the opportunity to discuss any issues during your visit to the Allergy clinic.



North West Paediatric Allergy Network

Subcutaneous Immunotherapy (SCIT)

Information for parents and carers

www.allergynorthwest.nhs.uk

Allergic rhinitis (hay fever) is one of the most common allergies, affecting 1 in 4 young people in Britain. In a small proportion of patients, maximum treatment (with antihistamines, nasal steroids and eye drops) is not effective. Some of these patients may benefit from immunotherapy.

What is immunotherapy?

Allergies occur when the body reacts to foreign proteins (allergens). In allergic rhinitis, nose symptoms (runny, itchy or blocked nose) or eye symptoms (itchy watery eyes) occur after contact with these allergens e.g. grass or tree pollen, house dust mite, dogs and cats. In children, grass and tree pollen allergy is an important cause of hay fever.

Immunotherapy is the only treatment that can change the underlying allergic disease. The alternative is to continue with medicines that dampen down the symptoms such as antihistamines and steroids.

How is immunotherapy given?

Immunotherapy is given by either injection under the skin (subcutaneous immunotherapy [SCIT]), or as a drop under the tongue (sublingual immunotherapy [SLIT]). Giving increasing amounts of the allergen trains the immune system to ignore the allergen (immune tolerance).

Subcutaneous immunotherapy (SCIT) is used by our allergy service, particularly for grass pollen and tree pollen allergies. In a few children it is also used to treat children with bee or wasp venom allergies.

What are the benefits of immunotherapy?

Nose and eye symptoms improve. Patient's often feel better, are less sleeping and are able to concentrate more on their school work and outside activities.

Allergen immunotherapy may also reduce the likelihood of getting asthma.

Worm M, et al, Randomized controlled trials define shape of dose response for Pollinex Quattro. . .Allergy, 2018
Rosewich M, et al. Pollinex Quattro: an innovative four injections immunotherapy in allergic rhinitis. Hum Vaccin Immunother, 2013

What is Pollinex Quattro SCIT?

Pollinex Quattro SCIT is a grass and tree pollen therapy used for children with very troublesome hay fever. Four injections are given under the skin. They are usually in the upper arm. Injections have to be given each week for one month on three consecutive years.

The vaccine course is usually given in winter, when the pollen count is at its lowest.

How safe is SCIT?

SCIT is safe. Over 500,000 doses of Pollinex Quattro have been injected worldwide. The medicine is well tolerated in the vast majority of patients.

The most common side effects are itchiness and swelling in the arm. A dose of antihistamine and/or paracetamol will usually reduce these symptoms.

More serious reactions are uncommon, but include a widespread rash, wheeze or a drop-in blood pressure (feeling faint), which is why the treatment is always given in hospital. These reactions almost always occur soon after the injection and can be identified and effectively treated while the patient is still in hospital. This is why we ask you to stay for one hour after the injection.

You will need to be well on the day of injection. The injection should be postponed if you feel unwell with a cold or flu, or you have, or are getting over, an asthma attack.

What happens if I miss a dose?

If you are not able to have one of your injections, it will be given the following week.

Produced January 2019, to review 2022
SCIT - North West Paediatric Allergy Implementation Group
This information should be used in conjunction with advice from a medical professional